











<p>Mushroom/Toadstool-shaped: No Gills</p>	<p>dry scaly cap and teeth = <i>Sarcodon</i> dry cap and pores = <i>Boletus/Boletellus/Stobilomyces</i> slimy cap and pores = <i>Suillus</i></p>
<p>Underside of cap with downward-oriented spines or “teeth”; Leathery cap 2 to 6 inches across, light to dark brown, depressed in center and scaly; Grows on ground</p> <p>Scaly Tooth—<i>Sarcodon imbricatus</i></p>	 <p>George Barron, University of Guelph, Canada</p>
<p>Cap dry and with large, dense, dark, soft warts, 2.5 to 5 inches across; Flesh white, changing to red and then blackish when cut; Pores start white, becoming blackish; Stalk 2.5-4 inches long, covered to some extent with same warts as cap.</p> <p>Old Man of the Woods—<i>Strobilomyces sp.</i></p>	 <p>Brian Shelton, University of Guelph, Canada</p>
<p>Cap 3-6 inches across, surface dry, gray, rather thick; Flesh gray-white or dull yellow, becoming pinkish when cut after a while; Pores white to gray, bruise brown; Stalk 2-4.5 inches long, whitish to yellowish, with darker netting, base often crooked</p> <p>Gray Bolete—<i>Boletus griseus</i></p>	 <p>George Barron, University of Guelph, Canada</p>
<p>pores bruise blue or gray</p> <p>Cap 2-6 inches across, surface dry, purplish to dark red at first, becoming yellowish-mottled; Flesh yellow, often becoming more golden when cut; Pores yellow to brownish, bruise blue; Stalk 2.5-4 inches long, also bruises blue.</p> <p>Red-and-Yellow Bolete—<i>Boletus bicolor</i></p>	 <p>John Walker, University of Vermont</p>

pores bruise same color as flesh	<p>Cap 1-2 inches across, surface dry, brownish yellow, with erect fiber-clusters; Flesh yellow, unchanged when cut; Pores cream-colored, turning to greenish-yellow, does not bruise; Stalk 3-4 inches long, tapering towards cap and curving at base, netted by flesh-colored or red ridges</p> <p>Russell's Bolete—<i>Boletellus russellii</i></p>	 <p>George Barron, U. of Guelph, Canada</p>
	<p>Cap 1-2.5 inches across, surface dry, leathery, becoming cracked yellowish to olive; Flesh white, becoming yellow when cut; Pore surface does not bruise; Stalk 1-2 inches, same color as cap, with dark flecks;</p> <p>Parasitic on puffballs Parasitic Bolete—<i>Boletus parasiticus</i></p>	 <p>George Barron, University of Guelph, Canada</p>
	<p>Cap 1-4.5 inches across, slightly slimy or dry, red with pattern of yellowish cracks; Flesh creamy-yellow, slowly becoming reddish when cut; Pores yellow, aging to brown, does not bruise; Stalk 1.5-3 inches long, colored like the cap.</p> <p>Painted Bolete—<i>Suillus pictus</i></p>	 <p>P. Super, NPS</p>
	<p>Cap 1.5-4 inches across, slimy, surface chrome yellow, tinged or spotted reddish; Flesh yellowish, turning pinkish-brown then yellow when cut; Smells slightly sour; Stalk 1-2 inches long, yellow with reddish-brown glandular dots.</p> <p>Slippery Jack or Chicken Fat Mushroom <i>Suillus americanus</i></p>	 <p>George Barron, University of Guelph, Canada</p>
	<p>Cap 1.5-4 inches across, slimy, brownish to reddish-chestnut with dark stains; Flesh white, does not change color when cut; does not bruise; Stalk short, 0.5-1.5 inches, whitish or yellow and tapering downward.</p> <p>Short-stalked Bolete—<i>Suillus brevipes</i></p>	 <p>George Barron, University of Guelph, Canada</p>
	<p>Cap 1-3 inches across, slimy, orange to yellow, with dense tufts of fibers; Flesh pale yellow, unchanged if cut, but can stain paper greenish-yellow; Pores pale yellow to brownish, does not bruise; Stalk 1.5 to 2.5 inches long, yellowish, speckled with reddish dots</p> <p>Hairy Bolete—<i>Suillus hirtellus</i></p>	 <p>R. Vilgalys, Duke University</p>